

Cell Boost 7a and 7b supplements

HYCLONE MEDIA AND SUPPLEMENTS

HyClone™ Cell Boost™ 7a and Cell Boost 7b supplements are intended for use in combination with HyClone ActiPro™ and ActiSM™ media as part of the ActiPro cell culture system formulated to provide high yields of recombinant proteins in processes when using Chinese hamster ovary (CHO) cell lines. They are available as a convenient ready-to-use liquid feed solution or as a dry powder concentrated format to be hydrated before use (Fig 1). Cell Boost 7a and 7b are chemically defined (CD), animal-derived component-free (ADCF), and optimized for high-yield protein production in fed-batch processes. Cell Boost 7a and 7b do not contain any growth factors (such as insulin), peptides, hydrolysates, phenol red, or 2-mercaptoethanol, ensuring batch-to-batch consistency and increased cell culture process efficiency.

The ActiPro system of media and supplements is designed with adaptation and production in mind. The ActiPro system is suitable for general biomanufacturing with CHO cell lines such as CHO-GS, CHO-K1, CHO-DG44, and CHO-S.

Features and benefits

- Demonstrated high yields of recombinant proteins (e.g., mAbs > 5 g/L) in fed-batch processes.
- Chemically defined ADCF formulations.
- Concentrated feeds that complement each other and are used together.
- Accompanied with detailed protocols for use to maximize output.

Cell Boost supplements are used as an essential part of your cell culture feed strategy to enhance cell culture performance and increase product yield. Cell Boost 7a and 7b are intended for use together in defined concentrations.

The supplements should be added to the cultivation vessel as individual solutions and should not be mixed in advance as this will cause precipitation. The recommended ratio of Cell Boost 7a to 7b is 10 to 1 (v/v) (example feeding strategy Cell Boost 7a at 3% and Cell Boost 7b at 0.3%). The total amount of feed added and the specific feeding regime will need to be adjusted according to the nutritional requirements of each specific clone. The supplements are available as a powder or in a ready-to-use liquid format for research and process development. The liquid feed supplements meet testing specifications and are stable up to 12 months in PETE bottles and 9 months in bioprocessing bags at 2°C to 8°C (Fig 2).



Fig 1. ActiPro, ActiSM, and Cell Boost 7a and 7b are designed for high yields of recombinant proteins in fed-batch culture processes with CHO cells.




Fig 2. Stability study testing of the Cell Boost 7a (A) and 7b (B) liquid feed supplements was performed using CHO-S cell line to determine viable cell density (VCD) compared with control (freshly hydrated powder). The study was conducted using three separate batches of Cell Boost 7a and 7b liquid feed supplements stored at 2°C to 8°C and used in fed-batch cultures of CHO-S cells 0, 3, 6, 9, and 12 months after preparation.

Specifications

Cell Boost 7a and 7b specifications for liquid feed solution and dry powder formats:

- Protein-free
- Growth factor-free
- Phenol red-free
- 2-mercaptoethanol-free
- Hydrolysate-free
- Chemically defined and ADCF formulated

Cell Boost 7a

Cell Boost 7a has a pH close to neutral and contains amino acids, vitamins, salts, trace elements, and glucose.

Cell Boost 7b

Cell Boost 7b has an alkaline pH and is a concentrated solution of amino acids.

Safety declaration

Cell Boost 7b is a hazardous chemical:

- Causes skin irritation
- Causes severe eye irritation
- Can cause respiratory irritation

Please take appropriate safety precautions prior to using Cell Boost 7b.

1. Wear protective gloves
2. Wear eye or face protection
3. Avoid breathing dust

If inhaled, move the individual to fresh air and keep at rest in a position comfortable for breathing. Call a poison center or physician if you feel unwell.

Store Cell Boost 7b in a locked location.

Dispose contents and container in accordance with all local, regional, national, and international regulations.

Shelf-life

Cell Boost 7a and 7b liquid feed supplements stored in PETE bottles have a shelf-life up to 12 months when stored at 2°C to 8°C.

Cell Boost 7a and 7b liquid feed supplements stored in bioprocess bags have a shelf-life up to 9 months when stored at 2°C to 8°C.

Please refer to the product label for the expiry date of your shipment.

Storage

Store Cell Boost 7a and 7b liquid feed supplement at 2°C to 8°C, protected from light. In addition, powder supplement should be stored in a dry environment, protected from light and stored at 2°C to 8°C.

Quality control testing for liquids

Quality control test specifications for Cell Boost 7a and 7b liquid formats are listed in Table 1.

Table 1. Test specifications*

	Cell Boost 7a	Cell Boost 7b
Appearance	Clear solution	Clear solution
Endotoxin (EU/mL)	≤ 1 EU/mL	≤ 1 EU/mL
Sterility	No growth	No growth
pH	6.45–6.80	11.0–11.4
Osmolality (mOsmol/kg)	247–303 (diluted 1:5)	218–266 (diluted 1:5)
HPLC amino acid profile	Complies with profile	Complies with profile

*Refer to certificate of analysis for actual results.

Suggested preparation for powder supplement

To ensure proper reconstitution, please follow the protocol for your Cell Boost supplement.

Required equipment and materials include:

- a. Mixing vessel
- b. Stirrer
Note: a magnetic stirrer can be used for small-scale reconstitutions up to 5 L. An overhead or bottom-mounted impeller is recommended for larger volumes.
- c. Calibrated pH meter
- d. Calibrated osmometer
- e. Highly purified water such as water for injection (WFI)

Reconstitution of Cell Boost 7a powder supplement

1. Fill a clean mixing vessel up to 80% of the final volume with high-quality purified water, such as WFI, at ambient temperature (18°C to 25°C) and start stirring. For example, start with 700 to 800 mL of water for a preparation of 1 L Cell Boost 7a solution.

Note: the stirring should be vigorous enough to quickly draw the powder below the liquid surface, but not too fast to cause formation of air bubbles and excessive foaming.

Note: Adjust the initial volume based on the concentration of NaOH used for pH adjustment (see step 3).

2. Add 181.04 g/L Cell Boost 7a powder slowly to the vessel, avoiding formation of clumps. Mix for 30 minutes. The solution will remain cloudy in this step but should be clear of any clumps or dry powder residues. The color will be bright red.
3. Slowly add 18.6 mL/L of a 5 N NaOH solution or 9.3 mL/L of a 10 N NaOH solution and mix for 60 minutes. After this step, the solution will be clear.
4. Adjust the pH to between 6.6 and 6.8 by drop-wise addition of 5 or 10 N NaOH or HCl. After adjusting, continue stirring for an additional 60 min to ensure that all components are completely dissolved.

Note: Our recommendation for pH adjustment of the dry powder medium (DPM) upon hydration is between 6.6 to 6.8 at 20°C ± 0.5°C. This is in our product specifications for release criteria of the DPM and is in the certificate of analysis (C of A) for this product. The pH specifications for powder hydration represent the optimal range for use at the time of hydration.

The pH for CB7a liquid tends to drift down with storage over time. The pH specifications for the liquid products are adjusted to account for this phenomenon. This is in our product specifications for release criteria of the liquid media (LM) and is in the C of A for this product in liquid form. In-process testing is generally done at a higher pH value of 6.7 at 20°C ± 0.5°C as target value. pH values down to 6.45 (Table 1) are not detrimental to the performance of the product.

- Adjust to the final volume with high-quality purified water, such as WFI, and stir for an additional 10 min.
- Measure and record the final pH and osmolality.
Expected values:
 - pH 6.45 to 6.80
 - Osmolality 247 to 303 mOsmol/kg (diluted 1:5)
- Sterilize immediately by membrane filtration. Use a low-binding filter membrane type, such as PVDF or PES. The supplement is a clear bright red liquid.
- Store the reconstituted supplement protected from light at 2°C to 8°C until use.

Reconstitution of Cell Boost 7b powder supplement

- Fill a clean mixing vessel up to 80% of the final volume with high-quality purified water, such as WFI, at ambient temperature (18°C to 25°C) and start stirring. For example, start with 700 to 800 mL of water for a preparation of 1 L Cell Boost 7b solution.

Note: the stirring should be vigorous enough to quickly draw the powder below the liquid surface, but not too fast to cause formation of air bubbles and excessive foaming.

Note: Adjust the initial volume based on the concentration of NaOH used for pH adjustment (see step 3).

- Add 94.6 g/L Cell Boost 7b powder slowly to the vessel, avoiding formation of clumps. Mix for 30 minutes. The solution will remain cloudy in this step but should be clear of any clumps or dry powder residues.
- Slowly add 160.5 mL/L of a 5 N NaOH solution or 80.25 mL/L of a 10 N NaOH solution and mix for 60 minutes. After this step, the solution will be clear.
- Adjust the pH to between 11.0 and 11.4 by drop-wise addition of 5 or 10 N NaOH or HCl. After adjusting, continue stirring for an additional 60 min to ensure that all components are completely dissolved.

Note: the pH will gradually decrease with longer mixing times. Use caution when adjusting pH. Over adjusting can cause the osmolality to be out of specification.

- Adjust to the final volume with high-quality purified water, such as WFI, and stir for an additional 10 min.
- Measure and record the final pH and osmolality.
Expected values:
 - pH 11.0 to 11.4
 - Osmolality 218 to 266 mOsmol/kg (diluted 1:5)
- Sterilize immediately by membrane filtration. Use a low-binding filter membrane type, such as PVDF or PES. The supplement is a clear colorless or slightly yellow liquid. The color will turn darker yellow to brown over time.
- Store the reconstituted supplement protected from light at 2°C to 8°C until use.

Preparation note

Cell Boost 7a and 7b powder supplements can be conveniently reconstituted using single-use mixers, such as the Xcellerex™ XDM mixers (Fig 3).

Product handling

Store powder supplement protected from moisture in a tightly sealed container.

Store the liquid feed supplement at 2°C to 8°C, away from light.

General culture recommendations

Cell Boost 7a and 7b are used as an essential part of your cell culture feed strategy and should be used together in defined concentrations to enhance cell culture performance and increase product yield.

- Cultures should be incubated at 37°C in a 7.5% CO₂ environment.
- Maintain adapted cells by establishing a mid-logarithmic growth phase subculturing schedule.
- Suggested seeding density of cultures 3.0×10^5 cells/mL; viability should be > 90%.

Process conditions

ActiPro medium and Cell Boost 7a and 7b supplements are recommended for use in a CO₂ atmosphere. Equilibration of ActiPro media in 7.5% CO₂ will result in a starting pH of 7.15 ± 0.05 . During the cultivation, pH control can be managed by varying the CO₂ concentration or by addition of base such as NaHCO₃ or NaOH. The culture temperature should be adjusted according to the requirements of the specific clone or target product. ActiPro medium and Cell Boost 7a and 7b supplements have demonstrated excellent results both under constant temperature conditions and in biphasic processes comprising a shift to a lower temperature.

Custom production

Formulations and delivery systems can be customized to your specific process requirements or optimized to maximize process yields.

Rapid Response Production (RRP)

Our RRP program manufactures up to 200 L of your custom prototype formulation within seven working days of your request. Use our RRP service to expedite the development and testing of custom media for your biopharmaceutical manufacturing process.



Fig 3. XDM Quad Mixing System, with a powerful motor and magnetically locked impeller, effectively mixes even highly viscous materials.

Related products

Product specifications for Cell Boost 7a and 7b supplements and related products are listed in Table 2.

Table 2. Product specifications for ActiPro media and supplements

Specification	ActiSM	ActiPro	Cell Boost 7a	Cell Boost 7b
L-glutamine	-	-	-	-
Glucose	Y	Y	Y	-
Phenol red	-	-	-	-
Proteins	-	-	-	-
Hydrolysates	-	-	-	-
2-mercaptoethanol	-	-	-	-
Poloxamer 188	Y	Y	Y	-

ActiSM medium

ActiSM is a lean, chemically defined ADCF medium that does not contain glycine, hypoxanthine, or thymidine. This medium was developed to be used as the first step in adapting cells to the ActiPro system of media and supplements. Once cells have undergone the adaptation process, we recommend using ActiPro medium for production.

ActiPro medium

ActiPro is a rich, chemically defined, and ADCF medium that does not contain hypoxanthine or thymidine and has been formulated for use in high-yield batch or fed-batch processes.

Technical support

Our cell culture medium specialists and technical support functions are happy to discuss your needs in getting the most out of your culture. In addition, we have an extensive service offering to help with, for example, process development, optimization, and scale-up. Please contact your local sales representative to learn more about the services we offer. To find a certificate or a MSDS for a specific product, please visit cytiva.com/certificates.

Ordering information

Product	Pack size	Format	Product code
HyClone Cell Boost 7a liquid supplement	500 mL bottle	Liquid	SH31119.01
	1000 mL bottle	Liquid	SH31119.02
	1 L bag*	Liquid	SH31119.03
	5 L bag*	Liquid	SH31119.04
	10 L bag*	Liquid	SH31119.05
	20 L bag*	Liquid	SH31119.06
	50 L bag*	Liquid	SH31119.07
	100 L bag*	Liquid	SH31119.08
	200 L bag*	Liquid	SH31119.09
HyClone Cell Boost 7a dry powder supplement	1 L bottle*	Powder	SH31026.07
	5 L bottle	Powder	SH31026.01
	10 L bottle*	Powder	SH31026.02
	25 L bottle*	Powder	SH31026.03
HyClone Cell Boost 7b liquid supplement	100 mL bottle	Liquid	SH31120.01
	500 mL bottle	Liquid	SH31120.02
	1000 mL bottle	Liquid	SH31120.03
	1 L bag*	Liquid	SH31120.04
	5 L bag*	Liquid	SH31120.05
	10 L bag*	Liquid	SH31120.06
	20 L bag*	Liquid	SH31120.07
	50 L bag*	Liquid	SH31120.08
	100 L bag*	Liquid	SH31120.09
	200 L bag*	Liquid	SH31120.10
HyClone Cell Boost 7b dry powder supplement	0.5 L bottle	Powder	SH31027.01
	1 L bottle*	Powder	SH31027.07
	5 L bottle*	Powder	SH31027.02
	10 L bottle*	Powder	SH31027.04

Related products	Pack size	Format	Product code
HyClone ActiPro powder medium	5 L	Powder	SH31037.01
	10 L*	Powder	SH31037.02
	25 L*	Powder	SH31037.05
HyClone ActiPro liquid medium	500 mL	Liquid	SH31039.01
	1000 mL	Liquid	SH31039.02
	1 L poly bag*	Liquid	SH31039.03
HyClone ActiSM powder medium	5 L	Powder	SH31038.01
	10 L*	Powder	SH31038.02
	25 L*	Powder	SH31038.05
HyClone ActiSM liquid medium	500 mL	Liquid	SH31040.01
	1000 mL	Liquid	SH31040.02
	1 L poly bag*	Liquid	SH31040.03
L-glutamine 200 mM	100 mL		SH30034.01
	500 mL		SH30034.02
	500 g*		SH30336.03

Note: powder product quantity is shown as the final volume of after powder reconstitution.

*Item is made to order. Lead times and minimum order quantities apply.

cytiva.com/hyclone

For local office contact information, visit cytiva.com/contact

Cytiva and the Drop logo are trademarks of Global Life Sciences IP Holdco LLC or an affiliate. ActiPro, ActiSM, Cell Boost, HyClone, and Xcellerex are trademarks of Global Life Sciences Solutions USA LLC or an affiliate doing business as Cytiva. All other third-party trademarks are the property of their respective owners.

© 2020–2021 Cytiva

All goods and services are sold subject to the terms and conditions of sale of the supplying company operating within the Cytiva business. A copy of those terms and conditions is available on request. Contact your local Cytiva representative for the most current information.

CY9773-27Jan21-DF

